

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup>:</b> C07H 17/00	A1	<b>(11) International Publication Number:</b> WO 94/26758 <b>(43) International Publication Date:</b> 24 November 1994 (24.11.94)
<b>(21) International Application Number:</b> PCT/US94/02547 <b>(22) International Filing Date:</b> 14 March 1994 (14.03.94) <b>(30) Priority Data:</b> 08/064,610 19 May 1993 (19.05.93) US <b>(60) Parent Application or Grant</b> (63) Related by Continuation US 08/064,610 (CON) Filed on 19 May 1993 (19.05.93) <b>(71) Applicant (for all designated States except US):</b> PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> YANG, Bingwei, V. [CN/US]; 27 Lincoln Road, Waterford, CT 06385 (US). <b>(74) Agents:</b> SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).		<b>(81) Designated States:</b> CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> INTERMEDIATE FOR AZITHROMYCIN  <b>(57) Abstract</b>  A compound having formula (III). Also, a process of making azithromycin comprising reducing the compound of formula (III) and N-methylating the reduced product. <div data-bbox="641 1129 1247 1501"><p style="text-align: right;">(III)</p></div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

INTERMEDIATE FOR AZITHROMYCINFIELD OF THE INVENTION


5 This invention relates to antibiotics, and particularly relates (1) to an intermediate per se, useful for making the known antibiotic azithromycin, (2) to a process for making azithromycin with the intermediate, and (3) to processes for making the intermediate.

10

BACKGROUND OF THE INVENTION

Erythromycin is an antibiotic formed during the culturing of a strain of *Streptomyces erythreus* in a suitable medium as taught in U.S. Patent No. 2,653,899. Erythromycin, which is produced in two forms, A and B, is represented by the following structure (I):

15



25

ERYTHROMYCIN

30


Erythromycin	R
A	-OH
B	-H

35

-2-

The structure reveals that the antibiotic is comprised of three main portions: a sugar fragment known as cladinose, a second sugar moiety containing a basic amino substituent known as desosamine and a fourteen membered lactone ring referred to as erythronolide A or B or as the macrolide ring.

Azithromycin is the U.S.A.N. (generic name) for 9a-aza-9a-methyl-9-deoxo-9a-homoerythromycin A, a broad spectrum antibacterial compound derived from erythromycin A. Azithromycin was independently discovered by Bright, U.S. Pat. No. 4,474,768 and Kobrehel et al., U.S. Pat. No. 4,517,359, and was named N-methyl-11-aza-10-deoxo-10-dihydro-erythromycin A in these patents. It has the following structure (II) wherein the numbering system conventionally employed is shown:



30 The above patents also disclose that (II) and certain derivatives thereof possess antibacterial properties.


35

-3-

In particular, the procedure for making azithromycin from erythromycin A involves relatively strong reaction conditions as described, for example, in Djokic et al.,  
5 *J. Chem. Soc. Perkin Trans. I*, 1881, (1986), wherein the preparation of 10-dihydro-10-deoxo-11-azaerythromycin A from an imino ether precursor was effected by catalytic hydrogenation in acetic acid under 70 atm of H<sub>2</sub>.

### SUMMARY OF THE INVENTION

10 In a first aspect, this invention provides a compound having formula III



20 (III)


Compound III, 9-deoxo-11-deoxy-9,11-epoxy-9,9a-didehydro-9a-aza-9a-homoerythromycin A, herein also referred to as a 9,11-imino ether, has utility as an  
25 intermediate for making azithromycin, and can be reduced to a direct precursor of azithromycin having formula IV, shown below. The precursor of Formula IV need only be N-methylated (position 9a) to produce azithromycin. Accordingly, in a further aspect, this invention provides a process of making azithromycin, comprising reducing the (9,11-imino ether) compound of formula III to 9a-aza-9-deoxo-9a-homoerythromycin  
30 A, a compound having formula IV

35

-4-

5

10




and thereafter N-methylating the said compound of formula IV.

In a further aspect, this invention provides a process of making an intermediate of formula III, comprising isomerizing a compound of formula V, 9-deoxy-6-deoxy-6,9-epoxy-9,9a-didehydro-9a-aza-homoerythromycin A:

20

25




in a suitable solvent.

30

In still a further aspect, this invention provides an additional process of making an intermediate of formula III, comprising treating a compound of formula VI

35

-5-




with tosyl chloride and pyridine in diethyl ether at a temperature of less than about 10°C for a time between about 0.5 and about 50 hours.

#### DETAILED DESCRIPTION

A general reaction scheme for (1) making the intermediate 9,11-imino ether and (2) using the intermediate 9,11-imino ether to make azithromycin is shown in Scheme 1:


Scheme I






Scheme I, Con't.

STEP B




Reduction

STEP C



formic acid  
formaldehyde

(II)



In the above scheme the starting material in step A, compound (VI) is the E-isomer of 9-deoxo-9-hydroxyimino erythromycin A and can be made by known procedures such as the straightforward reaction of Erythromycin A with hydroxylamine to produce the erythromycin E-oxime as the major isomer. The oxime is treated with tosyl chloride in the presence of pyridine as a base and in a suitable solvent such as a dialkyl ether (e.g. diethyl ether), thereby initially forming a corresponding O-tosyl oxime which is a suitable precursor in the Beckmann rearrangement of ketoximes. Importantly, the temperature should be maintained below -10°C to favor formation of the 9,11 imino ether, compound III, over compound V, the 6,9 imino ether, and compound VII. Generally it is preferred to maintain the said temperature below -20°C, and most preferred to maintain the temperature below -40 °C. At the low temperatures employed, and depending on the exact temperature employed, the reaction can take as long as several days to go to completion, although appreciable amounts of 9,11-imino ether III are available from workup of the reaction medium after much shorter reaction times, e.g., on the order of a day. The amount of tosyl chloride employed is at least equivalent to the amount of erythromycin A 9-E-oxime. To ensure completion of the reaction within a reasonable time the tosyl chloride can be used in excess, with a 2:1 equivalents ratio being preferred. The mixture will typically contain a mixture of compounds V, III, and VII, per Scheme I, Step A, which can be conventionally resolved by column chromatography separation, typically employing silica gel having a particle size of 230-400 mesh ASTM (commercially available, for example, as Silica Gel 60 from EM Science, Gibbstown, NJ) with toluene, chloroform and triethylamine mixed in a ratio, respectively, of 20:1:1 as the eluting solvent system.

The fact that 9,11-imino ether III is made per Step A is surprising in view of literature precedent (Djokic et al., supra) wherein the erythromycin A 9-E-oxime of erythromycin A yielded 6,9-imino ether V only.

If appreciable amounts of 6,9-imino ether V are formed, it can be isomerized to compound III by dissolving it in a suitable solvent. Any of a number of solvents can be employed such as tetrahydrofuran, (THF), lower alcohols, (e.g. methanol, ethanol, or propanol) and halogenated hydrocarbons, with chlorinated hydrocarbons such as chloroform and methylene chloride being preferred. Deuterated analogs of the chlorinated solvents (e.g. deuteriochloroform) can also be employed. To speed the rate of isomerization a catalytic amount of acid can be added to the reaction medium. The

acid employed is not critical so long as it is not used in an amount which results in cleavage of either of the sugar fragments from the macrolide ring. Organic acids (e.g. trifluoroacetic acid, p-toluenesulfonic acid) or inorganic (hydrochloric, sulfuric) acids may be employed. It is preferred to use camphorsulphonic acid in an amount of 0.1 equivalent per equivalent of compound V at a temperature of from 15°C to reflux, typically room temperature, for a period sufficient for isomerization to occur, typically a period of from about 12 hours to about 7 days or longer. Conversions of essentially 100% after about 7 days in deuteriochloroform are feasible using this novel procedure. Total yields of the intermediate III of over 90%, based on the weight of the starting material, are facily obtainable by combining compound III obtained directly from scheme I with compound III obtained by isomerizing compound V (also obtained from scheme I) in deuteriochloroform.

Compound III can be reduced to compound IV, the direct precursor of azithromycin, by any of a number of conventional methods, but advantageously under much milder conditions than those known in the art for reducing macrolide imino ethers. Reduction can be effected under a hydrogen pressure of about 50 psi in the presence of platinum dioxide catalyst and in glacial acetic acid solvent. Yields approaching 90% are obtainable within reaction times of 48 hrs. Other methods of reduction employing conventional reducing agents such as sodium borohydride are also feasible. Reactions employing borohydride are typically conducted with stirring in a suitable solvent such as methanol at a temperature of 0° to room temperature, and employing at least one equivalent of borohydride. Workup can proceed as in Djokic et al., supra.

Compound IV can be methylated to obtain azithromycin as conventionally known in the art, for example by the Eschweiler-Clark reaction in which compound IV is reacted with a combination of formic acid and formaldehyde, most suitably performed with a 1-3 molar excess of formaldehyde and formic acid in an appropriate solvent, preferably in a halogenated hydrocarbon, e.g., chloroform or carbon tetrachloride. The reaction is generally conducted at reflux for a period of 2 to 8 hours. The azithromycin can be isolated by conventional means such as by simple solvent evaporation. If further purification is desired, such can be effected conventionally, for example column chromatography through silica gel employing an eluting solvent comprising, by volume, 3-10% chloroform, and 0.1-1% ammonium hydroxide.

The invention will now be illustrated by means of the following examples which are not, however, to be taken as limiting:

5

Example 19-Deoxo-11-deoxy-9,11-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound III)

**METHOD A:** (Step A in Scheme I): Erythromycin A 9-oxime (6.11 g, 8.16 mmol) was dissolved in pyridine (45 ml) and cooled to -45°C. To this solution was added a  
10 precooled (-45°C) solution of *p*-toluenesulfonyl chloride (3.2 g, 16.9 mmol) and pyridine (10 ml) in ether (25 ml). The reaction mixture was stirred at -45°C for six hours and stood at -20°C for 12 hours. The mixture was poured into cold water and stirred while the pH was adjusted to 5 with aqueous 2M HCl. The solution was extracted with 75 mL  
15 quantities of methylene chloride twice. After separation, the aqueous layer was extracted with 75 mL of chloroform at pH 7 and 9, respectively (pH adjusted with saturated aqueous solution of potassium carbonate). Each extract was separately washed with brine and dried over magnesium sulfate. The extract from pH 7, upon evaporation *in vacuo*, gave a slightly yellow solid, containing a mixture of 9-deoxo-11-deoxy-9,11-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound III) and 9-deoxo-6-deoxy-6,9-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound V)  
20 at 1.2/1 ration (4.69g, 6.42 mmol, yield: 78.6%). The pH 9 extract was evaporated to yield a white solid containing 9a-aza-9a-homoerythromycin cyclic lactam (compounds VII) and 9-deoxo-6-deoxy-6,9-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (V) at  
25 2/1 ratio (0.49 g). A pure sample of III was obtained as follows: The mixture of V and III was dissolved in deuteriochloroform and stirred at room temperature for 12 hours. After removal of solvent *in vacuo*, the residue was chromatographed on silica gel (eluent: toluene/CHCl<sub>3</sub>/Et<sub>3</sub>N 20:1 :1) to afford the pure title compound as a white solid. <sup>13</sup>CNMR (CDCl<sub>3</sub>) 176.3, 102.2, 95.3, 83.0, 82.3, 81.1, 77.7, 76.7, 76.4, 75.1, 73.2, 72.7,  
30 70.7, 69.4, 65.9, 65.1, 63.5, 49.3, 43.4, 40.3, 39.6, 34.6, 34.6, 29.8, 28.7, 25.0, 24.2, 21.7, 21.66, 21.0, 19.3, 18.1, 17.4, 11.3, 10.9.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, partial): 5.03 (br d, 1H), 4.81 (dd, J=10.1, 1.8 Hz, 1h), 4.61(d, J=7.3 Hz, 1h), 4.50 (D, J=4.7 Hz, 1H), 4.39 (d, J=8.5 Hz, 1H), 4.14 (m, 1H), 3.95 (m, 1H), 3.69 (d, J=5.9 Hz, 1H), 3.67 (s, 1H), 3.54 (m, 1H), 3.31 (s, 3H), 3.24 (m, 1H), 3.00 (t, J=9.7 Hz, 1H), 2.70 (m, 1H), 2.66 (m, 1H), 2.25 (s, 6H), 1.97 (s, 3H), 1.38 (d, J=6.6 Hz,

-11-

3H), 1.28 (s, 3H), 1.25 (d, J=6.4 Hz, 3H), 1.22 (s, 3H), 1.21 (s, 3H), 1.20 (s, 3H), 1.18 (s, 3H), 1.175 (s, 3H), 1.14 (d, J=7.2 Hz, 3H), 0.88 (t, J=7.2 Hz, 3H).

- 5 FAB-mass: m/e: 731, 573, 398, 158; high resolution mass calcd for  $C_{37}H_{67}N_2$ : 731.4696; found 731.4744.

Method B: A solution of 9-deoxo-6-deoxy-6,9-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound V) (100 mg, 3.57 mmol) in  $CDCl_3$  (1.5 ml) was stirred at room temperature for 36 hours.  $^1H$  NMR spectrum of an aliquot sample indicated  
10 that the reaction mixture contained 9-deoxo-6-deoxy-6,9-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound V) and 9-deoxo-11-deoxy-9,11-deoxy-9,11-epoxy-9,9a-dihydro-9a-aza-9a-homoerythromycin A (compound III) at ratio 1.7/1. After seven days, the isomerization of compound V to III was complete, as indicated by  $^1H$  NMR spectrum. Evaporation of the solvent afforded the title compound in quantitative yield;  
15 it was identical with that obtained according to method A.

### Example 2

#### 9-Deoxo-9a-aza-9a-homoerythromycin A (compound IV)

A solution of the title compound of example 1 (231 mg, 0.316 mmol) in acetic acid (20  
20 ml) was hydrogenated over  $PtO_2$  (13.1 mg, 0.058 mmol) under hydrogen (50 psi) at room temperature for 48 hours. The catalyst and the solvent were removed. The residue oil was dissolved in methylene chloride and washed with 10% aqueous potassium carbonate and brine, and dried over magnesium sulfate. The solvent evaporated *in vacuo* to afford the title compound pure (199 mg, 0.27 mmol, yield:  
25 85.8%); it was identical by comparison with a known sample of the title compound.

30


35

-12-


CLAIMS

What is claimed is:

- 5           1.     A compound having formula III



- 20           2.     A process of making azithromycin, comprising reducing a compound of  
formula III




- 35     to a compound of formula IV

-13-

5

10



15 and thereafter N-methylating the said compound of formula IV.


3. A process as defined in claim 2, wherein said compound of formula III is reduced with hydrogen in the presence of platinum dioxide catalyst.

4. A process as defined in claim 2, wherein said compound of formula IV is N-methylated by treating said compound of formula IV with a combination of formic  
20 acid and formaldehyde.

5. A method of making a compound having formula III

25

30




comprising treating a compound of formula VI

35

-14-

comprising treating a compound of formula VI




with tosyl chloride in the presence of pyridine at a temperature less than about -10°C.

6. A process as defined in claim 5; wherein said temperature is less than about -20°C.

7. A process as defined in claim 6, wherein said temperature is less than about -40°C.

8. A process of making a compound having formula III







## INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 94/02547

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 C07H17/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 508 795 (MERCK & CO. INC.) 14 October 1992 see page 12, line 45 - page 13, line 38 see page 26, line 42 - line 49; examples 9,10 ---	1-11
A	EP,A,0 137 132 (SOUR PLIVA FARMACEUTSKA) 17 April 1985 see claim 2 ---	1-11
A	J.CHEM. SOC. PERKIN TRANS. I 1986 pages 1881 - 90 S. DJOKIC 'Erythromycin Series. Part 11' cited in the application Scheme 2 see page 1187 - page 1189 --- -/--	1-11

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

13 September 1994

Date of mailing of the international search report

07.10.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 cpo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Bardili, W

# INTERNATIONAL SEARCH REPORT

In tional Application No  
PCT/US 94/02547

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>J. ORG. CHEM. vol. 57 , 1992 pages 4361 - 7 A.B. JONES 'New Macrolide Antibiotics' compound (19) see page 4364</p> <p style="text-align: center;">-----</p>	1-11

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/02547

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0508795	14-10-92	US-A- 5189159 CA-A- 2065674 JP-A- 5213964	23-02-93 11-10-92 24-08-93
-----			
EP-A-0137132	17-04-85	CA-A- 1231943 JP-C- 1350567 JP-A- 60038393 JP-B- 61014155 JP-A- 61148191	26-01-88 28-11-86 27-02-85 17-04-86 05-07-86
-----			